Open Access
J. Food Proc. & Nutr. Res., 2024, 1(1):43-49
ISSN (Online): 3078-9133; ISSN (Print):3078-9125
DOI: https://doi.org/10.22194/ JFPNR /24.1008
https://societyfia.org/journal/JFPNR

Optimizing Nutritional Biscuits: Enhancing Protein and Mineral Content for Improved Health Benefits

Nasir Shabbir[1*], Shabbir Ahmad[1], Muhammad Usman[2], Umar Farooq[2], Muhammad Sibt-eAbbas[1], Sadaf Yaqoob[3] and Shahrina Shakir[1]

[1]Department of Food Science and Technology, MNS-University of Agriculture, Multan, Pakistan; [2]Department of Human Nutrition and Dietetics, MNS-University of Agriculture, Multan, Pakistan; [3]Department of Food Science and Technology, Bahauddin Zakariya University, Multan, Pakistan
[*]Corresponding author's e-mail: nasirshabbir728@gmail.com

Malnutrition is an imbalance of energy and nutrients that is shown from physical status of children and adults. According to the WHO, more than 20 million children are severely malnourished under the age of five. In Pakistan, half of all children under the age of five are stunted and one in every ten infants is wasted. Malnutrition can be improved by fortification in food items with natural ingredients. Therefore, the present study was carried out to prepare nutritionally enriched biscuits by using wheat flour, chickpea, lentil and maize flour in different concentrations. The study was performed in the lab of Food Science & Technology, Volka Foods International (R & D lab) and Central Hi-Tech lab of MNS-University of Agriculture, Multan. The raw material was analyzed to determine macro and micro components. Afterwards, biscuits were prepared and analyzed for their nutritional composition like crude fiber, crude fat, crude protein, moisture, ash content and mineral (Fe & Zn) contents. "T5" (60% wheat+20% chickpea+20% maize) was best in terms of protein content, "T10" best in terms of fat content. Results showed that, hardness of biscuits decreased during storage days (1st, 7th and 14th days) while moisture content of biscuits increased with the time. Treatment "T9" color values was l* (63.85±0.937), a* (8.53±0.870) and b* (27.30±0.740) showed best results in terms of color analysis. According to the results, "T6 and T9" showed best results in terms of sensory characteristics like color, appearance, flavor, taste, aroma, texture and overall acceptability of biscuits. Concentration of 60% wheat flour, 20% chickpea flour and 20% lentil flour (T6) showed best results in terms of color, flavor and overall acceptability while concentration of 50% wheat flour, 20% chickpea flour, 20% maize flour and 10% lentil flour (T9) was best in terms of appearance, aroma, texture and taste in biscuits. Overall results showed that, T9 (Wheat flour 50%, Chickpea flour 20%, maize flour 20% and lentil flour 10%) had the best results compared to the other treatments. Sensory scores and moisture content both decreased significantly during the storage period.

Keywords: Malnutrition, composite flour, fortified biscuits, compositional analysis, sensory characteristics.

INTRODUCTION

Malnutrition is a condition when the body is deficient in the nutrients which require for development, maintenance and some tasks. Protein Energy Malnutrition (PEM) a body's requirements for protein, energy or both are not met (Sheetal *et al.*, 2013). Asian countries have severe malnutrition situation than other regions of the world, with 81.7 million stunted (55%), 33.8 million wasted (68.28%) and 18.8 million overweight (47%). Pakistan is unexpectedly one of the countries with the highest malnutrition rates, ranking 77 out of 113; it is also considered to have the worst stunting rates in the Southern Asian Countries (SAC). Stunting, wasting and underweight are more common in rural communities than in urban ones. In terms of gender-based statistics, males are more likely than females to be stunted, wasted, underweight and overweight. One out of every ten infants waste and more than 20 million children under the age of five suffer from severe malnutrition (Ahmad *et al.*, 2020).

Cereals are Gramineae grass family edible seeds or grains that are grown for their gastronomic attributes (Art of selecting, preparing, serving and enjoying fine food). Wheat, rice, rye, barley, oats, maize, sorghum, triticale and millet are among the grains farmed in many nations. The two most important crops in the world, accounting for more than half of grain production are wheat and rice (Mckevith, 2004). People have

Shabbir, N., S. Ahmad, M. Usman, U. Farooq, M. Sibt-eAbbas, S. Yaqoob and S. Shakir. 2024. Optimizing Nutritional Biscuits: Enhancing Protein and Mineral Content for Improved Health Benefits. Journal of Food Processing and Nutrition Research 1:43-49.
[Received 13 Jan 2024; Accepted 2 Apr 2024; Published 14 Nov 2024]
Attribution 4.0 International (CC BY 4.0)

been eating cereals for a long time and cereal grains are cultivated on large scale. They also contribute important part in the human nutrition as an energy source (Dias-Martins *et al.*, 2018). The FAO estimates that 2,706 million tonnes of cereal will be produced globally in 2019–2020, with a forecast of 2.741 million tonnes for the 2020–2021 season. Above 73% of the world's harvested land-dwelling is used to cultivate cereals which provide more than 60 percent of worldwide food consumption (Charalampopoulos *et al.*, 2002). Cereals contain nutrients such as carbohydrates, minerals, vitamins, proteins and fats. Some cereals also contain a lot of dietary fibre (insoluble and soluble). These grains' physiological properties support the prevention or control of diseases including hypertension, diabetes and heart risk (Baniwal *et al.*, 2021).

Wheat (*Triticum aestivum*) is the important cash crop for farmers and governments, as well as one of the world's most widely consumed cereals. It belongs to the *Triticum* genus (Tebben *et al.*, 2018). Wheat is still recognized as the "queen of cereals" due to its versatility in producing leavened foods. Wheat is the cereal that contributes in the majority of a person's daily diet (Serna-Saldivar, 2016). Most of the grain's protein and starch are found in the inner endosperm. It is kept apart from wheat that is being milled into flour. Protein, vitamins and phytic acid are more abundant in the aleurone and outer layers of the endosperm than they are in the inner endosperm (Kumar *et al.*, 2011). Wheat contributes 20% of the food's calories and approximately 55% of the carbohydrate. It has significant amounts of vitamins and minerals, including thiamine and vitamin B, as well as carbohydrates, protein, fat and micronutrients 78.10%, 14.70%, 2.10% and 2.10% respectively. About 72% of the protein in the seed is kept in the endosperm, which accounts for 8–15% of the total protein by grain weight (Kumar *et al.*, 2011). The content of protein in wheat ranges between 8.7 to 19% on a dry matter basis. Cereal proteins can be categorized according to their shape, biological role, solubility or chemical make-up. Albumins and globulins are soluble in water and salt respectively but both are insoluble in water. Gliadins (known as prolamins) are soluble in 70 to 90% ethanol and glutenins are insoluble in neutral aqueous, saline and alcohol solutions. While the previous two are primarily storage proteins, the first two are cytoplasmic or metabolically active proteins. A strong relationship exists between the total amount of storage proteins and the crude protein content of grains (Gutierrez-Alamo *et al.*, 2008).

Maize is main cereal crop that is harvest in many countries of the world and utilized as both human food and animal feed. Except in Antarctica, maize is the 3rd largest crop globally after wheat and rice. Most important for maize to thrive, it requires a specific environment, a certain amount of water and temperatures between 15 and 20 °C for germination (Ranum *et al.*, 2014). Starch, proteins and other compounds like the abundantly present anthocyanin and carotenoids which are often associated with pigmentation, make up the majority of the ingredients in maize grains. These substances are beneficial for human health (Kuhnen *et al.*, 2022). Maize grain is a tasty and nutritious part of the plant. It includes N-p-coumaryl tryptamine, N-ferrulyl tryptamine, vitamins (C, E, K, B1, B2, B3, B5 and B6), folic acid and selenium. Because it is lacking in the usual human diet, potassium, a primary nutrient present is significant (Kumar and Jhariya, 2013).

Pulses are members of the Leguminosae family which includes species that are eaten by people and domesticated animals. Pulses have a limited number of important amino acids like methionine, tryptophan and cysteine. They nonetheless have a protein content of between 21-25%. As compared to cereals, pulses have nearly double the amount of protein. Albumin and globulin were the two main fractions used to classify pulse proteins. The primary storage proteins in pulse seeds are globulins, which account for 35 to 72% of the total protein in the seeds. Albumins make up the majority of the remaining protein component. Pulses contain a wealth of vitamins and minerals (iron, zinc, calcium and magnesium). Cardiovascular diseases and an imbalance of the biological processes were caused by different minerals being deficient in different countries of the world (Singh, 2017). Due to their exceptional benefits, pulses play a vital role in addressing the triple burden of malnutrition. A wide variety of the population will gain nutritionally from the use of pulses which offer a promising alternative source of functional and nutritious proteins and are becoming more popular among consumers of cereals (Benali *et al.*, 2021).

Chickpea (*Cicer arietinum L.*) is most important legume with a lot of health and nutritional benefits. It is richest source of different phytochemicals like natural antioxidants such as flavonoid or phenolic compounds that reduce inflammation and oxidative stress associated with many chronic diseases and a rich source of non-digestible carbohydrates (Benali *et al.*, 2021). Chickpeas is a very healthy and cheap source of protein with a protein content around 24% that varies from 15% to 30% depending on various environmental factors. Chickpeas also have 60-65% carbs, 6%S fat and are vital source of B vitamins and minerals. Desi and Kabuli are the types which have several applications (Merga and Haji, 2019). The protein content of chickpeas is thought to be superior to that of other pulses, and they are a richest source of both protein and carbohydrates. Food based solutions try to enhance the quality of food and nutrition by improving the convenience and intake of variety of foods they targeted various nutritional deficits at the same time. It may be used to create nutritious value-added goods which can be used for low-income people as nutritious food in developing nations and patients suffering from lifestyle illnesses (Hirdyani, 2014).

Lentil (*Lens culinaris Medikus* subsp. *culinaris*) was cultivated in many countries around the world and has

become an essential food crop in the agriculture and food sector. Its seed is a richest source of protein, minerals contents and vitamins for human nutrition. Its ingestion with wheat or rice offers a balance of essential amino acids for human nutrition because of its maximum content of lysine and tryptophan (Sarker and Erskine, 2006). Lentils are exceptional source of protein, carbs, minerals and vitamins contents and play an important part in human nutrition. Compared to most other grain legumes, lentils cook more quickly and are easier to decorate because of their tiny size and flat form. This might explain why worldwide per capita consumption has increased over the previous 50 years (Sharpe *et al.*, 2013). A cup of lentils can provide 1.9 to 3.3 mg of zinc, 25 to 401 mg of selenium and 4.3 to 5.3 mg of iron (Johnson *et al.*, 2015).

MATERIALS AND METHODS

This study was conducted in the Food Science & Technology Department, MNS-University of Agriculture, Multan, Volka Foods International, and Central Hi-Tech Lab, Multan. The present study was carried out to prepare nutritionally enriched biscuits by using wheat flour, chickpea, lentil and maize flour in different concentrations. Grains (wheat, chickpea, maize, and lentil) were sourced from local markets in Multan.

***Milling and preparation of biscuits*:** Grains were cleaned to remove impurities (stones, dust, etc.), and then milled into whole meal flour using a micro mill (Culatti Type MFC) (Cappelli *et al*., 2020). Biscuit preparation followed the recipe of El-Gohery (2021) with modifications. Fat was melted and mixed with ground sugar for creaming. Flour mixtures (composite flour, skim milk powder, egg powder, and baking powder) were sifted and gradually added to the cream. Salt and a small amount of water were added to form dough, which was rolled, cut into shapes, and baked at 140–160°C for 20–25 minutes. Biscuits were cooled and stored in airtight packets.

***Proximate Analysis*:** Moisture content was determined using a moisture analyzer. Fat content was measured via Soxhlet extraction, while protein content was analyzed using the Kjeldahl method. Carbohydrate content (NFE %) was calculated by subtracting protein, fat, ash, and moisture from 100% (AACC, 2000).

***Mineral Analysis*:** Minerals (Fe and Zn) were analyzed using an atomic absorption spectrophotometer, following the (AOAC, 2006) wet digestion method. Samples were digested using a 7:3 HNO3:HCLO4 mixture, then diluted to 100 ml with distilled water.

***Color Analysis*:** Color was measured using a Hunter's colorimeter (Virginia, USA) to obtain parameters a* (redness), b* (yellowness), and l* (lightness) (Wani *et al*., 2015).



Figure 1. Flow chart of biscuits development.

Table 1. Treatment plan for biscuits development.

Treat.	Wheat flour (%)	Chickpea flour (%)	Maize flour (%)	Lentil flour (%)	Quinoa flour (%)
T_1	50	25	-	-	25
T_2	60	20	-	-	20
T_3	60	20	20	-	-
T_4	60	20	20	-	-
T_5	60	20	20	-	-
T_6	60	20	-	20	-
T_7	70	20	-	10	-
T_8	50	25	-	25	-
T_9	50	20	20	10	-
T_{10}	40	25	25	10	-

***Sensory Assessment*:** A panel of trained evaluators assessed the biscuits' flavor, texture, and overall acceptability on a scale of 1 to 9 (Lawless and Heymann, 2010).

***Statistical Analysis*:** Data were analyzed using one-way ANOVA and Tukey's test, with all measurements performed in triplicate. Statistical significance was determined at $p < 0.05$, using Statistics 8.1 software Steel and Torrie (1980).

RESULTS AND DISCUSSION

The current study was conducted in the Department of Food Science and Technology (FST) at MNS-University of Agriculture, Multan (MNS-UAM) and Volka Foods International. The planned research work was performed in Nutrient analytical lab. In current study, biscuits were analyzed for the proximate composition and other analysis. The obtained results in the research work are discussed below.

***Proximate Analysis of Biscuits*:** The proximate composition of biscuits made from flours blends are shown in Table 1. The moisture content of samples was not significantly different from each other ($p>0.05$). The moisture content of biscuits ranged from 3.18% to 5.37% among the treatments. The results are resembled with the finding of Van Toan and Anh (2018) who reported 4.89% to 5.96% moisture content in biscuits. The protein contents was not significantly different ($P>0.05$). The protein content of biscuits ranged from 6.75% to 9.34% among the treatments. Protein content of biscuits are found similar to the result of Pasqualone *et al.* (2020) who reported 5.06% to 10.66% of protein content in biscuits. Ash content showed highly significant difference. The ash content of biscuits ranged from 3.79% to 9.45% among the treatments. Fibre content is varies non-significant between different treatments. Mean values regarding the Fat content results ranged from 13.68% to 37.74% among the treatments. Results for fiber content of biscuits found in the current study are similar to the result of Van Toan and Anh (2018), who reported 0.33% to 0.91% of fibre content in biscuits. NFE content ranged from 40.96% to 65.21% among the treatments. NFE percentage of biscuits are found resembled to the findings of Farzana and Mohajan (2015) who reported 56.38% to 65.62% of NFE content of biscuits.

***Mineral Analysis*:** Iron content of biscuits is high as compared with the study of Hemalatha *et al.* (2007), who reported 6.64 to 7.83 mg/100g in biscuits sample. Zinc content of biscuits are similar with the study of Doner and Ege (2004), who reported 0.58 to 2.50 mg/100g in biscuits sample.

***Color Analysis*:** Mean values of color of biscuits are given in Table 3. Parameter l* results were noticed in the range of 42.42±0.869 to 66.68±0.872 and parameter a* results were observed in the range of 5.34±0.897 to 9.28±0.767. Maximum reading was 30.44±0.956% and minimum reading was 15.20±0.934 in parameter b*. Recent results of color analysis of biscuits are the resemblance with the findings of (Kumar *et al.*, 2015), who reported the 31.64 to 35.39 b* value, 10.66 to 13.53 a* value and 55.00 to 67.95 L* parameter of color of crust in biscuits, respectively. When

***Texture Analysis*:** The current findings regarding texture of biscuits are similar with the finding of Tyagi *et al.* (2007), Laguna *et al.* (2012) and Kumar *et al.* (2015), who reported 4.46N to 19.00N, 2.7N to 42.60N and 13.25N to 28.68N respectively of texture of biscuits.

***Sensory Analysis*:** This parameter describes the success or

Table 2. Mean values of proximate analysis.

Treatments	Protein%	Ash%	Fat%	Fiber%	Moisture%	NFE%	Iron (mg/100g)	Zinc (mg/100g
T1	7.60±0.71	3.79±0.83	29.04±0.99	0.52±0.16	3.45±0.59	55.23±0.50	21.4±0.05	0.13±0.04
T2	7.30±0.82	4.08±0.22	29.41±0.95	0.83±0.15	3.18±0.50	54.15±0.34	19.6±0.08	0.37±0.04
T3	7.38±0.82	3.87±0.81	26.35±0.76	0.59±0.10	4.26±0.62	57.26±0.52	10.8±0.09	0.11±0.17
T4	6.75±0.93	7.64±0.70	23.73±0.70	0.48±0.25	4.62±0.85	56.61±0.56	8.3±0.04	0.10±0.08
T5	9.34±0.95	7.11±0.96	28.65±0.84	0.43±0.23	5.37±0.71	49.03±0.52	14.6±0.04	0.07±0.05
T6	8.50±0.99	7.81±0.66	28.04±0.45	0.57±0.17	4.06±0.96	48.93±0.76	8.5±0.04	0.04±0.05
T7	8.58±0.97	8.23±0.68	25.84±0.79	0.51±0.05	3.33±0.80	53.46±0.57	16.5±0.20	0.02±0.05
T8	9.14±0.99	9.45±0.53	25.12±0.57	0.44±0.07	3.45±0.63	51.49±0.91	12.7±0.06	0.05±0.02
T9	7.44±0.95	8.86±0.95	13.68±0.72	0.53±0.04	3.59±0.51	65.21±0.31	23.0±0.17	0.08±0.14
T10	7.48±0.96	7.49±0.89	37.74±0.71	0.49±0.08	4.85±0.96	40.96±0.29	12.6±0.04	0.05±0.20

Table 3. Mean values of texture and color analysis.

Treatments	Hardness	l*	a*	b*
T1	19.43±0.122	42.42±0.869	5.34±0.897	15.20±0.934
T2	18.48±0.537	58.75±0.826	7.37±0.557	23.38±0.766
T3	18.18±0.282	66.68±0.872	7.58±0.666	26.74±0.905
T4	16.89±0.420	65.43±0.938	7.83±0.938	26.33±0.950
T5	15.22±0.525	59.68±0.957	8.56±0.712	27.32±0.639
T6	16.33±0.592	60.37±0.756	8.43±0.643	22.29±0.842
T7	20.77±1.642	58.54±0.528	9.28±0.767	26.05±0.583
T8	17.77±0.555	62.17±0.651	7.42±0.736	25.26±0.799
T9	14.60±1.148	63.85±0.937	8.53±0.870	27.30±0.740
T10	16.74±0.680	64.66±0.978	6.69±0.623	30.44±0.956

Table 4. Mean values of sensory evaluation.

Treatments	Color	Appearance	Texture	Aroma	Flavor	Taste	Overall Acceptability
T1	4.73±0.703	5.46±0.516	5.67±0.488	5.80±0.561	4.60±0.507	4.40±0.507	4.67±0.617
T2	4.27±0.458	4.47±0.516	4.47±0.516	4.40±0.507	4.53±0.516	4.40±0.507	4.20±0.414
T3	6.60±0.507	6.27±0.458	6.27±0.458	5.20±0.414	5.60±0.507	5.47±0.516	5.80±0.414
T4	6.73±0.458	6.80±0.414	6.40±0.507	5.87±0.352	6.60±0.632	6.33±0.488	6.60±0.507
T5	6.87±0.352	6.67±0.488	6.73±0.458	6.13±0.516	5.80±0.414	5.93±0.594	6.87±0.352
T6	7.67±0.488	7.20±0.414	7.27±0.458	7.40±0.508	7.47±0.516	7.27±0.458	8.07±0.594
T7	4.80±0.414	4.33±0.488	4.73±0.458	5.00±0.378	4.73±0.458	4.40±0.507	5.00±0.535
T8	5.87±0.743	7.00±0.926	5.73±0.458	6.27±0.458	6.40±0.508	5.87±0.352	6.13±0.352
T9	7.67±0.488	7.27±0.458	7.73±0.458	7.47±0.516	6.93±0.458	7.87±0.352	7.87±0.352
T10	6.93±0.594	7.73±0.458	5.80±0.704	6.87±0.352	6.87±0.352	7.13±0.352	7.00±0.000

acceptance of a food product among the client or consumer. The findings of treatments indicates that the highest value was 8.07±0.516 in T6, while 4.20±0.414 was lowest value in T2. Overall acceptability of biscuits are similar with the finding of Karki (2016) and Chandra *et al.* (2015) who reported 5.83 to 7.06 and 8.14 to 8.60 respectively of overall acceptability of biscuits sample.

***Conclusion*:** Malnutrition remains a serious public health concern around the world, particularly in poor nations, due to insufficient food quantity and quality. Bakery products can play a crucial role in overcoming malnutrition in children by providing essential nutrients, improving dietary diversity, and enhancing food accessibility. Compared to other snack foods, biscuits have a larger consumer base, a reasonably long shelf life and decent flavor. Protein and micronutrients (Fe and Zn) are necessary substances that people need in various amounts throughout their lives to coordinate a variety of physiological processes for maintaining health. According to the compositional analysis of nutritionally enriched biscuits, it can be concluded that T5 have higher protein content, T10 have higher fat content, T6 have higher ash content, T2 have higher fiber content and T9 have higher NFE content. According to the results, T9 showed best results in terms of color of biscuits. According to the results, T6 and T9 showed best results in terms of color, texture, flavor, appearance, aroma, taste and overall acceptability of biscuits among all samples.

***Authors contributions statement*:** Nasir Shabbir and Shabbir Ahmad designed, completed the experiments; Muhammad Usman, Sadaf Yaqoob and Shahrina Shakir prepared the draft; Dr. Sibt-e-Abbas, Umar Farooq reviewed and finalized the draft.

***Informed consent*:** N/A.

***Conflict of interest*:** The authors declare no conflict of interest.

REFERENCES

AACC. 2000. Approved methods of the American Association of Cereal Chemists. 11th Ed. AACC Press, Inc. St. Paul, Minnesota, USA.

Ahmad, D., M. Afzal and A. Imtiaz. 2020. Effect of socioeconomic factors on malnutrition among children in Pakistan. Future Business Journal 6:1-11.

AOAC (Association of Official Analytical Chemists). 2006. Official Methods of Analysis. The Association of Official Analytical Chemists, 18th Edition. Washington, DS. USA.

Baniwal, P., R. Mehra, N. Kumar, S. Sharma and S. Kumar. 2021. Cereals: Functional constituents and its health benefits. Journal of Pharmaceutical Innovation 10:343-349.

Benali, A., Y. En-Nahli, Y. Noutfia, A. Elbaouchi, M.R. Kabbour, F. Gaboun, E.H. El Maadoudi, N. Benbrahim, M. Taghouti and M. Ouhssine. 2021. Nutritional and technological optimization of wheat-chickpea-milk powder composite flour and its impact on rheological and sensorial properties of leavened flat bread. Foods 10:18-43.

Cappelli, A., N. Oliva and E. Cini. 2020. Stone milling versus roller milling: a systematic review of the effects on wheat flour quality, dough rheology, and bread characteristics. Trends in Food Science and Technology 97:147-155.

Chandra, S., S. Singh and D. Kumari. 2015. Evaluation of functional properties of composite flours and sensorial attributes of composite flour biscuits. Journal of Food Science and Technology 52:3681-3688.

Charalampopoulos, D., R. Wang, S. Pandiella and C. Webb. 2002. Application of cereals and cereal components in functional foods: a review. International Journal of Food Microbiology 79:131-141.

Dias-Martins, A.M., K.L.F. Pessanha, S. Pacheco, J.a.S. Rodrigues and C.W.P. Carvalho. 2018. Potential use of pearl millet (*Pennisetum glaucum (L.) R. Br.*) in Brazil: Food security, processing, health benefits and nutritional

products. Food Research International 109:175-186.

Doner, G. and A. Ege. 2004. Evaluation of digestion procedures for the determination of iron and zinc in biscuits by flame atomic absorption spectrometry. Analytica Chimica Acta 520:217-222.

El-Gohery, S.S. 2021. Effect of different treatments on nutritional value of lima bean and its utilization in biscuit manufacture. Food Science and Nutrition 12:372-391.

Farzana, T. and S. Mohajan. 2015. Effect of incorporation of soy flour to wheat flour on nutritional and sensory quality of biscuits fortified with mushroom. Food Science and Nutrition 3:363-369.

Gutierrez-Alamo, A., P.P. De Ayala, M. Verstegen, L. Den Hartog and M. Villamide. 2008. Variability in wheat: factors affecting its nutritional value. Journal of World's Poultry Science 64:20-39.

Hemalatha, S., K. Platel and K. Srinivasan. 2007. Zinc and iron contents and their bioaccessibility in cereals and pulses consumed in India. Food Chemistry 102:1328-1336.

Hirdyani, H. 2014. Nutritional composition of Chickpea (*Cicerarietinum-L*) and value added products: A review. Indian Journal of Community Health 26:102-106.

Johnson, C.R., P. Thavarajah, S. Payne, J. Moore and J.B. Ohm. 2015. Processing, cooking, and cooling affect prebiotic concentrations in lentil (Lens culinaris Medikus). Journal of Food Composition and Analysis 38:106-111.

Karki, R. 2016. Study on the preparation of biscuit and cake from amaranthus and sorghum. Journal of food science and technology Nepal 9:79-84.

Kuhnen, S., D.T. Holz, J.R. Moacyr, I.N. Piccinin and L.C. Pinheiro Machado Filho. 2022. Effect of pasture management on bioactive compounds of Lolium multiflorum and Avena strigosa for dairy cows and its effect on milk quality. Agroecology and Sustainable Food Systems 46:3-22.

Kumar, D. and A.N. Jhariya. 2013. Nutritional, medicinal and economical importance of corn: A mini review. Research Journal of Pharmaceutical Sciences 2:7-8.

Kumar, K.A., G. Sharma, M. Khan, T. Govindaraj and A. Semwal. 2015. Development of multigrain premixes—its effect on rheological, textural and micro-structural characteristics of dough and quality of biscuits. Journal of Food Science and Technology 52:7759-7770.

Kumar, P., R. Yadava, B. Gollen, S. Kumar, R.K. Verma and S. Yadav. 2011. Nutritional contents and medicinal properties of wheat: a review. Journal of Life Sciences Research. 22:1-10.

Laguna, L., P. Varela, A. Salvador, T. Sanz and S.M. Fiszman. 2012. Balancing texture and other sensory features in reduced fat short-dough biscuits. Journal of Texture Studies 43:235-245.

Lawless, H.T. and H. Heymann 2010. Sensory evaluation of food: principles and practices, Springer.

Mckevith, B. 2004. Nutritional aspects of cereals. Nutrition Bulletin 29:111-142.

Merga, B. and J. Haji. 2019. Economic importance of chickpea: Production, value and world trade. Cogent food and agriculture 5:1615718.

Pasqualone, A., B. Laddomada, F. Boukid, D.D. Angelis and C. Summo. 2020. Use of almond skins to improve nutritional and functional properties of biscuits: An example of upcycling. Foods 9:1705-1718.

Ranum, P., J.P. Pena-Rosas and M.N. Garcia-Casal. 2014. Global maize production, utilization and consumption. Annals of the New York Academy of Sciences 1312:105-112.

Sarker, A. and W. Erskine. 2006. Recent progress in the ancient lentil. Journal of Agricultural Science 144:19-29.

Serna-Saldivar, S.O. 2016. Cereal grains: Properties, processing and nutritional attributes, Boca Raton, CRC press, Mexico.

Sharpe, A.G., L. Ramsay, L.-A. Sanderson, M.J. Fedoruk, W.E. Clarke, R. Li, S. Kagale, P. Vijayan, A. Vandenberg and K.E. Bett. 2013. Ancient orphan crop joins modern era: Gene-based SNP discovery and mapping in lentil. BMC Genomics 14:1-13.

Sheetal, A., V.K. Hiremath, A.G. Patil, S. Sajjansetty and S.R. Kumar. 2013. Malnutrition and its oral outcome: A review. Journal of Clinical and Diagnostic Research 7:178-182.

Singh, N. 2017. Pulses: an overview. The Journal of Food Science and Technology. 54:853-857.

Steel, R.G.D. and J.H. Torrie 1980. Principles and procedures of statistics, a biometrical approach, McGraw-Hill Kogakusha, Ltd.

Tebben, L., Y. Shen and Y. Li. 2018. Improvers and functional ingredients in whole wheat bread: A review of their effects on dough properties and bread quality. Trends in Food Science and Technology 81:10-24.

Tyagi, S., M. Manikantan, H.S. Oberoi and G. Kaur. 2007. Effect of mustard flour incorporation on nutritional, textural and organoleptic characteristics of biscuits. Journal of Food Engineering 80:1043-1050.

Van Toan, N. and V.Q. Anh. 2018. Preparation and improved quality production of flour and the made biscuits from purple sweet potato. Journal of Food and Nutrition Research 4:1-14.

Wani, S.H., A. Gull, F. Allaie and T.A. Safapuri. 2015. Effects of incorporation of whey protein concentrate on physicochemical, texture, and microbial evaluation of developed cookies. Cogent Food and Agriculture 1:102-111.